

FIRST COHOMOLOGY, RIGIDITY AND DEFORMATIONS OF ISOMETRIC GROUP ACTIONS

DAVID FISHER

ABSTRACT. In 1964, Weil gave a criterion for local rigidity of a homomorphism from a finitely generated group Γ to a finite dimensional Lie group G in terms of cohomology of Γ with coefficients in the Lie algebra of G . This note announces a generalization of Weil's result to a class of homomorphisms into certain infinite dimensional Lie groups, namely diffeomorphism groups of compact manifolds. This gives a criterion for local rigidity of group actions which implies local rigidity of: (1) all isometric actions of groups with property (T), (2) all isometric actions of irreducible lattices in products of simple Lie groups and (3) a certain class of isometric actions of a certain class of cocompact lattices in $SU(1, n)$.

1. A COHOMOLOGICAL CRITERION FOR LOCAL RIGIDITY AND APPLICATIONS

In 1964, André Weil showed that a homomorphism π from a finitely generated group Γ to a Lie group G is *locally rigid* whenever $H^1(\Gamma, \mathfrak{g}) = 0$. Here π is locally rigid if any nearby homomorphism is conjugate to π by a small element of G , $\mathfrak{g}$ is the Lie algebra of G , and Γ acts on $\mathfrak{g}$ by the composition of π and the adjoint representation of G . Weil's proof also applies to G an algebraic group over a local field, but in all cases, his use of the implicit function theorem forced G to be finite dimensional. This note announces the following generalization of Weil's theorem to some cases where G is an infinite dimensional Lie group.

Theorem 1.1. *Let Γ be a finitely presented group, (M, g) a compact Riemannian manifold and $\pi : \Gamma \rightarrow \text{Isom}(M, g) \subset \text{Diff}^\infty(M)$ a homomorphism. If $H^1(\Gamma, \text{Vect}^\infty(M)) = 0$, the homomorphism π is locally rigid as a homomorphism into $\text{Diff}^\infty(M)$.*

The proof of Theorem 1.1 generalizes Weil's argument, with Hamilton's implicit function theorem in place of the standard implicit function theorem [Ha1, Ha2]. To apply Hamilton's theorem, I work with a tame exponential map from $\text{Vect}^\infty(M)$ to $\text{Diff}^\infty(M)$ which is a local diffeomorphism. This exponential map is defined by taking a vector field V to the map $\text{Exp}(V)(x) = \exp_x V_x$ where $\exp_x : TM_x \rightarrow M$ is the Riemannian exponential map. It is important not to use the map defined by flowing to time

Author partially supported by NSF grants DMS-0226121 and DMS-0541917.

one along V , as this has bad geometric properties, e.g. [Ha1, I.5.5.2]. Much of the proof of Theorem 1.1 does not require that the original action preserve a metric, see Theorem 2.1 below. That result is more difficult than Weil's, since we need to work with a more delicate implicit function theorem and since our exponential map is not equivariant. The main technical difficulty in proving Theorem 1.1 from Theorem 2.1 is producing the tame splittings required by Hamilton's theorem. Tameness is an assumption on Frechet manifolds and maps between them that is necessary for Hamilton's Nash-Moser implicit function theorem [Ha1, Ha2]. See section 2 for a more detailed discussion of the proof. Complete proofs of this result and the applications we now discuss will appear in [F1].

One can interpret Theorem 1.1 more dynamically. The homomorphism π defines an action of Γ on M by isometries and we will abuse notation by using π for both the action and the homomorphism. Theorem 1.1 says that any C^∞ action π' of Γ which is C^∞ close to π (which is equivalent to a homomorphism $\pi' : \Gamma \rightarrow \text{Diff}^\infty(M)$ which is close to π) is conjugate to ρ by a small C^∞ diffeomorphism. In e.g. [FM1, FM2], this condition is called $C^{\infty,\infty}$ *local rigidity* of π . I will use this dynamical language when discussing applications. The question of whether one could prove local rigidity of a group action by proving vanishing of $H^1(\Gamma, \text{Vect}(M))$ was first raised in [Z1], and more explicitly in the paragraph following the initial statement of Theorem 5.6 in [Z2]. This question has remained open, but many authors studied conditions under which $H^1(\Gamma, \text{Vect}(M))$ vanished, with a variety of assumptions on the regularity of the vector fields, see for example [H, Ko, L, LZ, Q, Z2]. Vanishing of $H^1(\Gamma, \text{Vect}(M))$ was labelled *infinitesimal rigidity* in the hope that vanishing of H^1 would imply local rigidity. Only the results in [LZ, Section 4] apply to isometric actions and so only those cohomology vanishing theorems yield applications of my results. In fact, I give a sharper version, Criterion 3.1, of Theorem [LZ, Theorem 4.1] in section 3 and will state below three applications of Theorem 1.1 and Criterion 3.1. In contexts similar to those considered by the authors listed above, one may be able to verify the hypothesis of Theorem 2.1 to prove local rigidity results, see [F1] for further discussion.

In earlier work Benveniste has proven some results on local rigidity of group actions, also by using Hamilton's implicit function theorem [Be]. Benveniste's results are special cases of Theorem 1.2 below. Benveniste uses the implicit function theorem to show that certain variations of geometric structures are trivial. These arguments are close in spirit to Weil's work in [We1, We2] and to the work of Calabi and Vesentini [C, CV]. My proof is based on a direct analysis of the space of actions as in Weil's work in [We3]. Kanai has also proven local rigidity results by considering variations of geometric structures, but without using Hamilton's theorem [Kan]. For a more detailed history of results on local rigidity see [F3].

I will now describe several applications of Theorem 1.2, all of which are proven by showing that $H^1(\Gamma, \text{Vect}^\infty(M)) = 0$. Using [LZ, Theorem 4.1], we obtain another proof of:

Theorem 1.2. [FM1] *Let Γ be a discrete group with property (T). Then any C^∞ , Riemannian isometric action of Γ on a compact manifold is $C^{\infty,\infty}$ locally rigid.*

Another, more novel, application is:

Theorem 1.3. *Let Γ be an irreducible lattice in a semisimple Lie group with real rank at least two. Then any C^∞ , Riemannian isometric action of Γ on a compact manifold is $C^{\infty,\infty}$ locally rigid.*

Theorem 1.3 applies to irreducible lattices in products of rank 1 groups. Not all of these groups have property (T) and so the theorem has many applications not covered by Theorem 1.2. For example it applies to actions of $SL(2, \mathbb{Z}[\sqrt{(2)}])$. In fact, Theorem 1.3 applies much more generally than stated. First, Γ can be an irreducible S -arithmetic lattice in a product of simple algebraic groups over different fields. Secondly, using a result from [Md], the result can be extended to apply irreducible lattices in fairly general products of locally compact topological groups.

For certain cocompact arithmetic lattices Γ in a simple group G , the arithmetic structure of Γ comes from a realization of Γ as the integer points in $G \times K$ where K is a compact Lie group. In this case it is always true that the projection to G is a lattice and the projection to K is dense. We say a Γ action is *arithmetic* if it is defined by projecting Γ to K , letting K act by C^∞ diffeomorphisms on a compact manifold M and restricting the K action to Γ . As a consequence of deep results of Clozel [Cl1, Theorem 3.2 and 3.5] concerning automorphic forms, Theorem 1.1, and Criterion 3.1, one has the following:

Theorem 1.4. *For certain principal congruence lattices $\Gamma < SU(1, n)$, any arithmetic action of Γ is $C^{\infty,\infty}$ locally rigid.*

Interestingly, some cocompact lattices in $SU(1, n)$ have homomorphisms ρ to $\mathbb{Z}$ [Ka, BW], and so have arithmetic actions with deformations provided the centralizer Z of K in $\text{Diff}^\infty(M)$ is non-trivial. Having centralizer allows one to deform the action along the image of the homomorphism $\rho \circ \sigma_t : F \rightarrow Z$ where $\sigma_t : \mathbb{Z} \rightarrow Z$ is any one parameter family of homomorphisms. This construction can also be applied to actions of lattices in $SO(1, n)$ where having a homomorphism to $\mathbb{Z}$ is much more common, see e.g. [Lu]. For Γ a lattice in $SU(1, n)$ I know of no example of a faithful Γ action with trivial centralizer which is not locally rigid.

On the other hand for lattices in $SO(1, n)$, even faithful, ergodic actions with trivial centralizer are not locally rigid. For a *standard lattice* Γ constructed using a quadratic form defined over a quadratic extension of $\mathbb{Q}$, one has a dense embedding of Γ in $SO(n+1)$.

Theorem 1.5. *For Γ a standard, arithmetic, cocompact lattice in $SO(1, n)$ as in the last paragraph, and ρ the Γ action on S^n defined by the dense embedding of Γ in $SO(n+1)$. Then the space of deformations of ρ is infinite dimensional.*

This construction uses a variant of Johnson-Millson bending and answers a question in [JM]. It will be discussed in detail in [F2].

2. ON THE PROOF OF THEOREM 1.1

This section outlines the proof of Theorem 1.1 without going through the background necessary to define a tame splitting or to state Hamilton's theorem. In the course of the outline, I state a more technical result for arbitrary actions, Theorem 2.1 below. The end of the section contains an informal discussion of tameness in the context of Theorem 1.1 and an indication of how it is checked.

Let M be a compact manifold and $\text{Vect}^\infty(M)$ the graded Frechet space of C^∞ vector fields on M . Given an affine connection ∇ there is an $\text{Aff}(M, \nabla)$ equivariant, tame exponential map Exp from $\text{Vect}^\infty(M)$ to $C^\infty(M, M)$ defined by $\text{Exp}(V)(x) = \exp_x(V_x)$ where $\exp_x$ is the exponential map defined by ∇ . The map Exp is a diffeomorphism from a neighborhood W of the zero vector field to a neighborhood U of the identity in $\text{Diff}^\infty(M) \subset \text{Vect}^\infty(M)$. Note that if ∇ is not complete, the map Exp may not be defined on all of $\text{Vect}^\infty(M)$, but this is not relevant to the arguments given here.

For the remainder of this section, fix a finitely presented group Γ and a presentation of Γ . This is a finite collection S of generators $\gamma_1, \dots, \gamma_k$ and finite collection of relators $w_1, \dots, w_r$ where each w_i is a finite word in the γ_j and their inverses. More formally, we can view each w_i as a word in an alphabet on k letters. Let $\pi : \Gamma \rightarrow \text{Diff}^\infty(M)$ be a homomorphism, which we can identify with a point in $\text{Diff}^\infty(M)^k$ by taking the images of the generators. We have a complex:

$$(1) \quad \text{Diff}^\infty(M) \xrightarrow{P} \text{Diff}^\infty(M)^k \xrightarrow{Q} \text{Diff}^\infty(M)^r$$

Where P is defined by taking ψ to $(\psi\pi(\gamma_1)\psi^{-1}, \dots, \psi\pi(\gamma_k)\psi^{-1})$ and Q is defined by viewing each w_i as a word in n letters and taking $(\psi_1, \dots, \psi_n)$ to $(w_1(\psi_1, \dots, \psi_n), \dots, w_k(\psi_1, \dots, \psi_n))$. Since $\text{Diff}^\infty(M)$ is proven to be a tame Frechet Lie group in [Ha1, Theorem II.2.3.5], it is easy to see that P and Q are smooth and tame. Letting Id be the identity map on M , it follows that $P(\text{Id}) = \pi$ and $Q(\pi) = (\text{Id}, \dots, \text{Id})$. Also note that $Q^{-1}(\text{Id}_M, \dots, \text{Id}_M)$ is exactly the space of Γ actions. Note that while this is a closed subset of $\text{Diff}^\infty(M)^k$, it is unclear that it is a manifold even in a neighborhood of any individual point. (In the finite dimensional setting, this set is an algebraic variety, and so is a manifold at “most” points.)

The tangent spaces of $\text{Diff}^\infty(M)$ at any point can be identified with $\text{Vect}^\infty(M)$. To avoid notational confusion, we let $A = \text{Diff}^\infty(M)$, $B = \text{Diff}^\infty(M)^k$ and $C = \text{Diff}^\infty(M)^r$. Then the complex in (1) becomes:

$$(2) \quad A \xrightarrow{P} B \xrightarrow{Q} C$$

and we can also consider the derivative complex of the complex in (2):

$$(3) \quad TA \xrightarrow{DP} TB \xrightarrow{DQ} TC$$

It is possible to compute the derivatives DP_{Id} and DQ_π explicitly where Id is the identity map on M , and see that the image of DP_{Id} is the space of coboundaries $B^1(\Gamma, \text{Vect}^\infty(M))$ and that the kernel of DQ_π is the space of cocycles $Z^1(\Gamma, \text{Vect}^\infty(M))$. Therefore the final hypothesis of Theorem 1.1 means that

$$\frac{\text{Ker}(DQ_\pi)}{\text{Im}(DP_{\text{Id}})} = 0.$$

If $\text{Diff}^\infty(M)$ were finite dimensional, then the implicit function theorem would immediately imply that in a neighborhood of π , the image of P maps onto the set of points in $\text{Diff}^\infty(M)^k$ where Q is equal to $(\text{Id}, \dots, \text{Id})$. In other words, in a neighborhood of π , the set of points which correspond to Γ actions is exactly equal to the set of points in the image of P , i.e. the conjugates of π . This would complete the proof and is more or less Weil's proof from [We3].

To apply Hamilton's implicit function theorem in this context, we need to find a tame splitting of equation (2), and in fact to find a smooth family of tame splittings over a neighborhood of U of Id_M in A . For a statement of Hamilton's theorem see [F1] or [Ha1, Theorem III.1.1.1]. The proof is [Ha2, Section 2]. More precisely, we need to find smooth tame maps VP and VQ that split the sequence

$$(4) \quad U \times \text{Vect}^\infty(M) \xrightarrow{DP} U \times \text{Vect}^\infty(M)^k \xrightarrow{DQ} U \times \text{Vect}^\infty(M)^r$$

i.e. tame smooth maps $VP : U \times \text{Vect}^\infty(M)^k \rightarrow U \times \text{Vect}^\infty(M)$ and $VQ : U \times \text{Vect}^\infty(M)^r \rightarrow U \times \text{Vect}^\infty(M)^k$ that are each linear in the second variable, such that $DP \circ VP + VQ \circ DQ$ is the identity on $\text{Vect}^\infty(M)^k$ for any $x \in U$. Note that the maps P and Q are $\text{Diff}^\infty(M)$ equivariant where $\text{Diff}^\infty(M)$ acts on A by translation and on B and C by conjugation. Since the $\text{Diff}^\infty(M)$ action on A is simply transitive, we can use this to reduce the problem of finding a splitting in a neighborhood U to the problem of finding a splitting at Id by [Be, Lemma 4.3]. Benveniste's lemma checks that the smooth, tame splitting at Id , when translated by the $\text{Diff}^\infty(M)$ action, provides a smooth tame splitting on all of U . It therefore suffices to find a splitting of the sequence:

$$(5) \quad \text{Vect}^\infty(M) \xrightarrow{DP_{\text{Id}}} \text{Vect}^\infty(M)^k \xrightarrow{DQ_\pi} \text{Vect}^\infty(M)^r$$

This yields the following:

Theorem 2.1. *Let Γ be a finitely presented group, M a compact manifold, and $\pi : \Gamma \rightarrow \text{Diff}^\infty(M)$ a homomorphism. If $H^1(\Gamma, \text{Vect}^\infty(M)) = 0$ and the sequence in line (5) admits a tame splitting, then the homomorphism π is locally rigid.*

See remarks in [F1] for some potential applications of Theorem 2.1. In the isometric case, we are assuming that $H^1(\Gamma, \text{Vect}^\infty(M)) = 0$, so that the map $DP_{\text{Id}} : \text{Im}(DP_{\text{Id}}) \rightarrow \text{Ker}(DQ_\pi)$ is surjective. Note that vanishing of H^1 does not necessarily imply that DP_{Id} has a tame inverse as there is no closed graph theorem in the tame Frechet category. To find the splitting at π and at $Q(\pi)$, we use the fact that $\text{Vect}^\infty(M)$ can be written as a (Hilbertian) direct sum of $\pi(\Gamma)$ invariant finite dimensional subspaces V_j , see the beginning of section 3 for more discussion. We can then split the complex in (4) by splitting each complex in the sequence of complexes:

$$(6) \quad V_j \xrightarrow{DP} V_j^k \xrightarrow{DQ} V_j^r.$$

Tameness can then be shown using the fact that the V_j are contained in eigenspaces of the Laplacian, the structure of isometric actions with vanishing cohomology and some estimates on “small divisors” that yield bounds on the maps used to split the complex in (6).

On tameness: The formal definition of tameness is quite involved, so we give an informal discussion in this setting. A Frechet space is a complete, Hausdorff, locally convex, topological vector space. The topology on a locally convex, topological vector space is always defined by a family of semi-norms. The Frechet space is called *graded* when we can choose an increasing family of semi-norms to define the topology. On $\text{Vect}^\infty(M)$, there is an increasing family of norms, the C^k norms $\|\cdot\|_k$ on vector fields, that define the topology. There is another increasing family of norms, the $W^{2,k}$ Sobolev norms $\|\cdot\|_{2,k}$ on vector fields. These two families of norms define the same topology on $\text{Vect}^\infty(M)$. More is true, if we normalize the metric so that M has volume 1, then the definitions of the norms combined with the Sobolev embedding theorems imply that:

$$(7) \quad \|v\|_{2,k} \leq \|v\|_k \leq C_k \|v\|_{2,k+r}$$

for any $v \in \text{Vect}^\infty(M)$ where $r = \frac{\dim(M)}{2}$ and C_k is a constant depending on k and the geometry of M . This statement is exactly the statement that these two families of norms on $\text{Vect}^\infty(M)$ are tamely equivalent in the language of [Ha1, Definition II.1.1.3].

More generally, given two graded Frechet spaces $(W, \|\cdot\|_j^W)$ and $(V, \|\cdot\|_l^V)$ and a linear map $L : W \rightarrow V$ is tame if there are positive

integers r and b such that:

$$(8) \quad \|Lv\|_j^V \leq C_j \|v\|_{j+r}^W$$

for every $v \in W$ and all $j \geq b$ and where C_j is a constant depending on j and L . This is [Ha1, Definition II.1.2.1].

The complex in line (4) can be split formally by splitting the complexes in line (6), but we need to see why this splitting is tame and why the maps constructed preserve smoothness. Since each V_j is contained in an eigenspace for some eigenvalue λ_j of the Laplacian on vector fields. Letting I be the identity on $\text{Vect}^\infty(M)$, and $\|\cdot\|_{2,k}$ the k -th Sobolev norm, we have:

$$(9) \quad \|v\|_{2,k} = \|(I + \Delta)^{k/2} v\|_{2,0} = (1 + \lambda_j)^{k/2} \|v\|_{2,0}$$

for every $v \in V_j$. The first equality is standard, and in fact is an equivalent definition for the Sobolev norm, and the second equality is obvious from the fact that $v \in V_j$.

This reduces the problem of splitting the complex in (2) to the problem of splitting the sequences in (6) by maps that are bounded by polynomials in λ_j . In the context of Theorem 1.1 the bounds are proven using arithmetic properties of $\pi(\Gamma)$ that we deduce from vanishing of $H^1(\Gamma, V_j)$ in a finite set of subspaces V_j . Tame estimates on the map VP are proven using the ideas discussed below in the context of verifying Criterion 3.1(2). Tame estimates on VQ are much more involved, see [F1].

3. PROVING $H^1(\Gamma, \text{Vect}^\infty(M)) = 0$

For applications of Theorem 1.1, I prove a criterion for the vanishing of $H^1(\Gamma, \text{Vect}^\infty(M))$. This criterion is a sharpening of [LZ, Theorem 4.1] and the proof is similar.

Using that eigenspaces of the Laplacian on vector fields are finite dimensional and $\text{Isom}(M, g)$ invariant, that $\text{Isom}(M, g)$ is compact and the Peter-Weyl theorem, we see that the space of vector fields splits as an infinite direct sum $\oplus_{j=1}^\infty V_j$ where each V_j is a finite dimensional, irreducible Γ -module, and contained in an eigenspace for the Laplacian. Let λ_j be the eigenvalue for the eigenspace containing V_j . This is a Hilbertian direct sum in either the L^2 topology or the Sobolev $W^{2,k}$ topology for any value of k . Since each V_j is contained in an eigenspace for the Laplacian on vector fields, it consists of C^∞ vector fields by a standard elliptic operator argument. Fix a finite generating set S for Γ and let $\|\cdot\|_2$ denote the L^2 norm on vector fields on M .

Criterion 3.1. *Let Γ be a finitely generated group, (M, g) a compact Riemannian manifold and $\pi : \Gamma \rightarrow \text{Isom}(M, g)$ a homomorphism. Then the following are sufficient conditions for vanishing of $H^1(\Gamma, \text{Vect}^\infty(M))$*

- (1) $H^1(\Gamma, V_j)$ vanishes for every j and

- (2) *there exist $\varepsilon > 0$ and a non-negative integer α such that for each V_j which is a non-trivial Γ -module and every $v_j \in V_j$ there is $\gamma \in S$ such that*

$$\|v - \pi(\gamma)v\|_2 \geq \varepsilon \lambda^{-\alpha} \|v\|_2.$$

Part (1) of the criterion provides the existence of a formal solution for any cohomological equation coming from $H^1(\Gamma, \text{Vect}^\infty(M))$. Part (2) guarantees a smooth solution of a smooth cohomological equation and provides tame estimates on the size of the solution in terms of the size of the coefficients in the equation. Given $v \in \text{Vect}^\infty(M)$, we can write $v = \sum_j v_j$ and then smoothness is equivalent to $\sum_j \lambda_j^n \|v_j\| < \infty$ for all positive integers n . Smoothness of solutions to smooth equations is this fact and condition (2) in Criterion 3.1.

For all of our applications, Criterion 3.1(2) can be verified by three methods. The first is to adapt the argument of the appendix of [Do] to our more general setting, see also [DoKr] for a related computation. The second is to use instead a deep arithmetic result of Clozel, which implies condition (2) with $\alpha = 0$, [Cl1]. More precisely, Clozel's result implies that all non-trivial V_j that can arise in this context are outside some neighborhood of the identity in the Fell topology on the unitary dual of Γ . This uses the fact that all representations of Γ which occur, when induced to the group G in which Γ is a lattice, occur in the automorphic spectrum of G . For the generalization of Theorem 1.3 for lattices in arbitrary products where Clozel's results do not apply, the reader should observe that any isometric action factors through a finite group, in which case verifying condition (2) is trivial. Lastly, we can show, as is done in [F1], that vanishing of certain $H^1(\Gamma, V_j)$'s implies certain arithmetic properties of $\pi(\Gamma)$. These properties are enough to show that each $\pi(\gamma)$ satisfies some estimate of diophantine type which is more than is needed to verify (2).

As remarked above, vanishing of $H^1(\Gamma, \text{Vect}^\infty(M))$ in the context of Theorem 1.2 was already observed in [LZ]. For Theorem 1.3, Criterion 3.1(1) is a consequence of Margulis's superrigidity theorem [Ma, Introduction, Theorem (3)]. For the more general lattices discussed in [Md] the vanishing of cohomology follows from the Monod's criterion for arithmeticity. More precisely, if the cohomology fails to vanish, Monod's results imply that Γ is S -arithmetic and then one can apply Margulis' result to obtain a contradiction. For Theorem 1.4, condition (1) of Criterion 3.1 is a special case of deep results of Clozel [Cl1, Theorem 3.2 and 3.5] concerning automorphic forms.

REFERENCES

- [Be] E.J.Benveniste, Rigidity of isometric lattice actions on compact Riemannian manifolds, *GAF* 10 (2000) 516-542.
- [BW] Borel, A.; Wallach, N. *Continuous cohomology, discrete subgroups, and representations of reductive groups*. Second edition. Mathematical Surveys and Monographs, 67. American Mathematical Society, Providence, RI, 2000.

- [C] E. Calabi On compact, Riemannian manifolds with constant curvature. I. 1961 *Proc. Sympos. Pure Math., Vol. III* pp. 155–180 American Mathematical Society, Providence, R.I.
- [CV] E. Calabi and E. Vesentini, On compact, locally symmetric Kähler manifolds. *Ann. of Math.* (2) 71 1960 472–507.
- [Cl1] Clozel, Laurent On the cohomology of Kottwitz’s arithmetic varieties. *Duke Math. J.* 72 (1993), no. 3, 757–795.
- [Cl2] Clozel, Laurent Demonstration de la conjecture τ . (French) [Proof of the τ -conjecture] *Invent. Math.* 151 (2003), no. 2, 297–328.
- [Do] Dolgopyat, Dmitry On mixing properties of compact group extensions of hyperbolic systems. *Israel J. Math.* 130 (2002), 157–205.
- [DoKr] D. Dolgopyat and R. Krikorian, On simultaneous linearization of diffeomorphisms of the sphere. *Duke Math. J.* 136 (2007), no. 3, 475–505.
- [F1] D. Fisher First cohomology and local rigidity of group actions, preprint.
- [F2] D. Fisher, Cohomology of arithmetic groups and bending group actions, in preparation.
- [F3] D. Fisher, Local rigidity of groups actions: past, present, future, in *Dynamics, Ergodic Theory and Geometry (Mathematical Sciences Research Institute Publications)*, 45–98, Cambridge University Press, 2007.
- [FM1] D. Fisher and G. A. Margulis, Almost isometric actions, property (T), and local rigidity. *Invent. Math.* 162 (2005), no. 1, 19–80.
- [FM2] D. Fisher and G. A. Margulis, Local rigidity for affine actions of higher rank Lie groups and their lattices, preprint, 59 pages, available at <http://mypage.iu.edu/~fisherdm/paperwork.html>, submitted.
- [Ha1] Hamilton, Richard S. The inverse function theorem of Nash and Moser. *Bull. Amer. Math. Soc. (N.S.)* 7 (1982), no. 1, 65–222.
- [Ha2] Hamilton, Richard S. Deformation of complex structures on manifolds with boundary. I. The stable case. *J. Differential Geometry* 12 (1977), no. 1, 1–45.
- [H] Hurder, Steven Infinitesimal rigidity for hyperbolic actions. *J. Differential Geom.* 41 (1995), no. 3, 515–527.
- [JM] Johnson, Dennis; Millson, John J. Deformation spaces associated to compact hyperbolic manifolds. *Discrete groups in geometry and analysis* (New Haven, Conn., 1984), 48–106, *Progr. Math.*, 67, Birkhäuser Boston, Boston, MA, 1987.
- [Kan] Kanai, M. A new approach to the rigidity of discrete group actions. *Geom. Funct. Anal.* 6 (1996), no. 6, 943–1056.
- [Ka] Kazhdan, David Some applications of the Weil representation. *J. Analyse Mat.* 32 (1977), 235–248.
- [Ko] Kononenko, A. Infinitesimal rigidity of boundary lattice actions. *Ergodic Theory Dynam. Systems* 19 (1999), no. 1, 35–60.
- [L] J. Lewis, Infinitesimal rigidity for the action of $SL_n(\mathbb{Z})$ on $\mathbb{T}^n$, *Trans. Amer. Math. Soc.* 324 no. 1 (1991) 421–445.
- [LZ] Lubotzky, Alexander; Zimmer, Robert J. Variants of Kazhdan’s property for subgroups of semisimple groups. *Israel J. Math.* 66 (1989), no. 1–3, 289–299.
- [Lu] Lubotzky, Alexander; Eigenvalues of the Laplacian, the first Betti number and the congruence subgroup problem. *Ann. of Math.* 144 (1996), no. 2, 441–452.
- [Ma] G. A. Margulis, *Discrete subgroups of semisimple Lie groups*, Springer-Verlag, New York, 1991.
- [Md] Monod, Nicolas Arithmeticity vs. nonlinearity for irreducible lattices. *Geom. Dedicata* 112 (2005), 225–237.
- [Q] Qian, Nantian Infinitesimal rigidity of higher rank lattice actions. *Comm. Anal. Geom.* 4 (1996), no. 3, 495–524.
- [We1] Weil, André On discrete subgroups of Lie groups. *Ann. of Math.* 72 (1960) 369–384.

- [We2] Weil, André On discrete subgroups of Lie groups II. *Ann. of Math.* 75 (1962) 578–602.
- [We3] Weil, André Remarks on the cohomology of groups. *Ann. of Math.* 80 (1964) 149–157.
- [Z1] R.J. Zimmer, Actions of semisimple groups and discrete subgroups, *Proc. Internat. Cong. Math.*, Berkeley, 1986, 1247-1258.
- [Z2] Zimmer, Robert J. Infinitesimal rigidity for smooth actions of discrete subgroups of Lie groups. *J. Differential Geom.* 31 (1990), no. 2, 301–322.

Department of Mathematics, Indiana University, Bloomington, IN 47401